


## MA PERINATAL HEPATITIS B PREVENTION PROGRAM

---

Massachusetts Department of Public Health  
Immunization Program  
MIAP 2016



1

---

---

---

---

---

---

---

---

### Presenter Disclosure Information

I, Theodora Wohler, have been asked to disclose any significant relationships with commercial entities that are either providing financial support for this program or whose products or services are mentioned during my presentations. I have no relationships to disclose.

I may discuss the use of vaccines in a manner not approved by the U.S. Food and Drug Administration

- But in accordance with ACIP recommendations

2

---

---

---

---

---

---

---

---

### Massachusetts Perinatal Hepatitis B Prevention Program

- **Program Mission:** Prevent perinatal and early childhood transmission of HBV through identification of Hepatitis B positive mothers, education and case management.
- **Program Goal:** To reduce infant HBV mortality and morbidity rates.

3

---

---

---

---

---


---

---

---

### Perinatal Hepatitis B and Hepatitis B Vaccine Requirements in MA

- 105 CMR 300.000: Requires all hepatitis B positive labs be reported to MDPH (1985)
- Perinatal Hepatitis B Program started (1989)
- Universal Hep B vaccination of all children (1991)
- Provides hepatitis B vaccine free of charge to birth hospitals and providers (1992)
- 105 CMR 130.627: Requires all pregnant women be tested for hepatitis B during each pregnancy (1993)



4

---

---

---

---

---

---

---

---

---

---

### Perinatal Transmission of Hep B

- An infant can acquire Hep B from:
  - An infected mother (transmitted at birth)
  - A chronically infected member of the household
- Risk of chronic infection
  - If mother is positive for HBsAg and HBeAg
    - 70% - 90% of infants infected
  - If mother positive for HBsAg only
    - 10% of infants infected
  - 90% of all those infected as infants become chronically infected

But, timely post-exposure prophylaxis protects up to 98% of infants (MMWR 10/9/15)

5

---

---

---

---

---

---

---

---

---

---

### Prevention of Perinatal Hepatitis B Virus Infection

- Begin treatment within 12 hours of birth
  - Hepatitis B vaccine (first dose) and HBIG (Hepatitis B Immunoglobulin)
- 2<sup>nd</sup> dose of Hep B vaccine: 1-2 months of age
- Final dose of Hep B Vaccine: 6 months of age
- Test for response (HBsAg and anti-HBs) after completion of at least 3 doses of the HepB series at 9 through 12 months of age (generally at the next well-child visit)

**Please note:**

- Pediarix dose given at 4 months of age is NOT the final dose. Another dose at 6 months of age is needed.
- Babies born to Hep B+ women that weigh less than 2000 grams at birth will need an extra dose of Hepatitis B vaccine.

6

---

---

---

---

---

---

---

---

---

---

**Hepatitis B Vaccine formulations to be used for pediatric vaccination**

Recombivax HB (Merck)

- Single antigen Hep B

Engerix-B (GSK)

- Single antigen Hep B

Pediarix (GSK)

- Combination vaccine
- DTaP, Hep B, IPV
- Approved for ages 6 weeks through 6 years old
- Approved for 3 doses at 2, 4, and 6 months
- Not approved for birth dose or booster doses
- Not recommended for use for revaccination
- May be used in infants whose mothers are HBsAg positive or status unknown

7

---

---

---

---

---

---

---

---

**Shortened Interval for Postvaccination Serologic Testing of Infants Born to Hepatitis B-Infected Mothers**

- Postvaccination serologic testing consists of two tests:
  - HBsAg (hepatitis B surface antigen)
  - Anti-HBs (antibody to hepatitis B surface antigen)
- CDC recently shortened the recommended age intervals for postvaccination serologic testing to **9-12 months of age**
  - In the past, recommendation was to perform test at age 9-18 months
- If completion of the series is delayed, the optimal timing for PVST to detect a vaccine response is 1-2 months after the last vaccine is administered.

<http://www.cdc.gov/mmwr/preview/mmwrhtml/mm6439a6.htm>

8

---

---

---

---

---

---

---

---

**Benefits and Goals of Shortened PVST Interval**

- Earlier identification of vaccine failures for referral
- Reduction in the time that non-responder infants are at risk for transmission from household contacts with Hep B
- Enables prompt revaccination for those infants needing a second series
- PVST occurring at an increasing interval after the final vaccine dose misclassifies some infants as vaccine non-responders and results in unnecessary revaccination

CDC. MMWR 2015;64:1118-1120.

9

---

---

---


---

---

---

---

---




---

---

---

---

---

---

---

---

---

---

---

---

### Interpretation of Hepatitis B Serologic Test Results

Hepatitis B serologic testing involves measurement of several hepatitis B virus (HBV)-specific antigens and antibodies. Different serologic "markers" or combinations of markers are used to identify different phases of HBV infection and to determine whether a patient has acute or chronic HBV infection, is immune to HBV as a result of prior infection or vaccination, or is susceptible to infection.

HBsAg	anti-HBc	anti-HBs	Interpretation
negative	negative	negative	Susceptible
negative	positive	negative	Immune due to natural infection
negative	negative	positive	Immune due to hepatitis B vaccination
positive	positive	negative	Acutely infected

• **Hepatitis B surface antigen (HBsAg):**  
A protein on the surface of hepatitis B virus; it can be detected in high levels in serum during acute or chronic hepatitis B virus infection. The presence of HBsAg indicates that the person is infectious. The body normally produces antibodies to HBsAg as part of the normal immune response to infection. HBsAg is the antigen used to make hepatitis B vaccine.

• **Hepatitis B surface antibody (anti-HBs):**  
The presence of anti-HBs is generally interpreted as indicating recovery and immunity from hepatitis B virus infection. Anti-HBs also develops in a person who has been successfully vaccinated.

<http://www.cdc.gov/hepatitis/hbv/pdfs/serologicchartv8.pdf>

11

---

---

---

---

---

---

---

---

---

---

---

---

### Screening for Risk Factors & Referrals for Perinatal HBV Infection

- Despite post exposure prophylaxis, mother to child transmission occurs in approximately 1% of infants born to HBsAg positive mothers.
- Risk factors for transmission include\*:
  - Hepatitis B e-antigen (HBeAg positive)
  - High HBV DNA concentration (DNA >20,000 IU/mL, approximately 112,000 viral copies per mL)
  - Elevated alanine aminotransferase (ALT) ( $\geq 19$  IU/L)
- Emerging evidence suggests that HBV treatment of pregnant women with antiviral agents in the 3<sup>rd</sup> trimester is safe and reduces rates of transmission
- CDC and ACOG have developed a *Screening and Referral Algorithm for HBV Infection among Pregnant Women*

\* Another risk factor is low birth weight (<2,000 grams) who receive the 1<sup>st</sup> dose of hepB vaccine before 1 month of age.

<http://www.cdc.gov/hepatitis/hbv/pdfs/prenatalhbsagtesting.pdf>

12

---

---

---

---

---

---

---


---

---

---

---

---




---

---

---

---

---

---

---

---

---

---

---

---

### Regional Immunization Nurses

Denise Dillon – Northeast	978-851-7261
Linda Jacobs – Southeast	508-984-0621
Lila Coverstone – Metro Boston	617-983-6811
Theodora Wohler – Metro West	617-983-6837
Katie Reilly, Nurse Manager	617-983-6833

14

---

---

---

---

---

---

---

---

---

---

---

---

# THANK YOU!



15

---

---

---

---

---

---

---

---

---

---

---

---