



DOSING FOR HEALTHCARE PROFESSIONALS

Use this chart to determine the proper dose of **TYLENOL**® for your patient. If possible, use weight to dose; otherwise, use age.[†]

[†] For pediatric products only

Recommended dose of **TYLENOL**® from your healthcare professional:

_____ dose of _____ product every _____ hours for _____ days

		Dosage						
Visit Tylenol.com for additional product information		Infants' TYLENOL® Oral Suspension*	Children's TYLENOL® Oral Suspension*	Children's TYLENOL® Chewables*	Children's TYLENOL® Dissolve Packs	TYLENOL® Regular Strength Tablets & Liquid Gels	TYLENOL® Extra Strength Caplets, Coated Tablets, Rapid Release Gels, & Dissolve Packs	TYLENOL® For Children + Adults Family*
mL = milliliter				Chew or crush tablets completely before swallowing				
Weight (lbs)	Age	Active ingredient: acetaminophen 160 mg (in each 5 mL)	Active ingredient: acetaminophen 160 mg (in each 5 mL)	Active ingredient: acetaminophen 160 mg (in each chewable tablet)	Active ingredient: acetaminophen 160 mg (in each powder)	Active ingredient: acetaminophen 325 mg (in each tablet/liquid gel)	Active ingredient: acetaminophen 500 mg (in each caplet /tablet /powder/gelcap)	Active ingredient: acetaminophen 160 mg (in each 5 mL)
6-11 lbs	0-3 mos	1.25 mL	The markings on the included dosing cup correspond to the doses listed below. This cup is designed for dosing children ages 2 - 11 years.	-	Do not use	OTC label does not include dosing for children under 6 years old	OTC label does not include dosing for children under 12 years old	The markings on the included dosing cup correspond to the doses listed below. This cup is designed for dosing ages 2 years and older.
12-17 lbs	4-11 mos	2.5 mL		-				
18-23 lbs	12-23 mos	3.75 mL		-				
24-35 lbs	2-3 yrs	5 mL	1 tablet					
36-47 lbs	4-5 yrs	The markings on the included syringe correspond to the doses listed above. This syringe is designed for dosing children ages 0 - 3 yrs.	7.5 mL	1½ tablets				
48-59 lbs	6-8 yrs		10 mL	2 tablets	2 powders	Children 6 years to under 12 years: Take 1 tablet/liquid gel every 4 to 6 hours while symptoms last. Do NOT take more than 5 liquid tablet/liquid gels in 24 hours	Repeat dose every 4 hours while symptoms last. Do NOT give more than 5 doses in 24 hours.	
60-71 lbs	9-10 yrs		12.5 mL	2½ tablets	2 powders			
72-95 lbs	11 yrs		15 mL	3 tablets	3 powders			
	Adults and children 12 years and over	-	-	-	-	Take 2 tablet/liquid gels every 4 to 6 hours while symptoms last. Do NOT take more than 10 tablet/liquid gels in 24 hours	Take 2 caplets/tablets/gelcaps/powders every 6 hours while symptoms last. Do NOT take more than 6 caplets/ tablets/gelcaps/ powders in 24 hours	20 mL

* The OTC label says 'ask a doctor' for dosing of children under 2 years. All Infants' TYLENOL® and Children's TYLENOL® products have the same strength of acetaminophen: 160 mg (in each 5 mL, tablet, or powder).

Remind parents & caregivers:

- Read & follow the label on all **TYLENOL**® products.
- Repeat dose **every 4 hours** while symptoms last.
- Do **NOT** give more than **5 doses in 24 hours**.
- Do **NOT** use with any other product containing acetaminophen.
- Use only the dosing device (syringe or dosing cup) that came with the product unless directed by a healthcare provider.

Remind patients:

- Read & follow the label on all **TYLENOL**® products.
- Do **NOT** use with any other product containing acetaminophen.
- Wait the right amount of time between doses



DOSING FOR HEALTHCARE PROFESSIONALS

Use this chart to determine the proper dose of **MOTRIN®** for your patient.
If possible, use weight to dose; otherwise, use age.†

† For pediatric products only

Recommended dose of **MOTRIN®**
from your healthcare professional:

_____ dose of _____ product every _____ hours for _____ days

Visit Motrin.com for additional product information		Dosage					
		Infants' MOTRIN® Concentrated Drops	Children's MOTRIN® Oral Suspension	Children's MOTRIN® Chewables	MOTRIN® IB Caplets	MOTRIN® IB Liquid Gels	MOTRIN® IB Migraine Liquid Gels
mL = milliliter							
Weight (lbs)	Age	Active ingredient: ibuprofen (NSAID)* 50 mg (in each 1.25 mL) <small>*Nonsteroidal anti-inflammatory drug.</small>	Active ingredient: ibuprofen (NSAID)* 100 mg (in each 5 mL) <small>*Nonsteroidal anti-inflammatory drug.</small>	Active ingredient: ibuprofen (NSAID)* 100 mg (in each chewable tablet) <small>*Nonsteroidal anti-inflammatory drug.</small>	Active ingredient: ibuprofen (NSAID)* 200 mg (in each caplet) <small>*Nonsteroidal anti-inflammatory drug.</small>	Active ingredient: ibuprofen (NSAID)* 200 mg (in each capsule) <small>*Nonsteroidal anti-inflammatory drug.</small>	Active ingredient: ibuprofen (NSAID)* 200 mg (in each capsule) <small>*Nonsteroidal anti-inflammatory drug.</small>
12-17 lbs	6-11 mos	1.25 mL	Infants' MOTRIN® and Children's MOTRIN® are different concentrations. The markings on the included dosing cup correspond to the OTC label doses listed below.	-	OTC label does not include dosing for children under 12 years old.	OTC label does not include dosing for children under 12 years old.	OTC label does not include dosing for children under 18 years old.
18-23 lbs	12-23 mos	1.875 mL		-			
24-35 lbs	2-3 years	Infants' MOTRIN® and Children's MOTRIN® are different concentrations. The OTC label does not include dosing directions for children ages 2 years and older.	5 mL	1 tablet			
36-47 lbs	4-5 years		7.5 mL	1½ tablets			
48-59 lbs	6-8 yrs		10 mL	2 tablets			
60-71 lbs	9-10 yrs		12.5 mL	2½ tablets			
72-95 lbs	11 yrs		15 mL	3 tablets			
-	-	If needed, repeat dose every 6-8 hours Do NOT use more than 4 times a day	If needed, repeat dose every 6-8 hours Do NOT use more than 4 times a day	If needed, repeat dose every 6-8 hours Do NOT use more than 4 times a day			

Remind parents & caregivers:

- Read & follow the label on all **MOTRIN®** products.
- If needed, repeat dose every 6-8 hours while symptoms last.
- Do **NOT** give more than 4 doses in 24 hours.
- Use only the dosing device (syringe or dosing cup) that came with the product.
- *Nonsteroidal anti-inflammatory drug

Use product only as directed.
© Johnson & Johnson Consumer Inc. 2023

Important Instructions For Proper Use:

- Always read and follow the label on all **MOTRIN®** products. Keep carton.
- Wait the right amount of time between doses
- If you have any questions, ask your healthcare professional which product may be right for you
- *Nonsteroidal anti-inflammatory drug